



## eVet Diagnostics

a Division of Veterinary Multi-Imaging, PLLC  
tel: 607 533 7512 fax: 607 533 7649

Page 1 of 2

*Report for case 392669*

Client:	
Organization:	Van Isle Veterinary Hospital
Email:	info@vanislevet.com
Phone / fax:	(250) 334-8400 / (250) 334-8403
Owner:	Claire Desrochers
Practitioner:	Dr. Yvette MacLean
Patient:	Bruno
Species/Breed:	Dog : Australian Labradoodle
Age:	6 Months
Gender:	male
Submitted Date:	8/21/2017 5:15:03PM
Completed Date:	8/29/2017 10:29:43AM

**Case status:** Completed

### Case notes:

Looking for a preliminary evaluation of the hips and elbows, future breeding stud

### The report:

392669 KW/lb

Four digitalized radiographic images containing projections of the pelvis and elbows were submitted for evaluation. The radiographs are dated 08/21/17.

#### Radiographic Findings:

Both femoral heads are located in their respective acetabula and are congruous with the acetabular rims.

Greater than 50% of the femoral heads are located medial to the dorsal acetabular rims.

No osteophyte formation is present on the femoral necks or acetabula.

Alignment of the patellae with the femurs is normal.

Muscle volume in both hind legs is within normal limits.

The anconeal process of both ulnae and proximocranial aspect of both radii are smooth with no evidence of osteophyte formation.

No sclerosis is associated with the ulnar notches.

Both medial coronoid processes are smooth in contour and a fragment associated with the medial coronoid processes is not identified.

**Radiologic Diagnosis:** Normal pelvis and elbows.

**Comments:** Radiographic changes to suggest elbow or hip dysplasia are not present. Based on the OFA scheme of evaluating the hips, the hips would be rated as good in conformation. Radiographs obtained at two years of age will need to be evaluated to completely exclude elbow and hip dysplasia and to establish a final rating on the coxofemoral joints.